arXiv:1001.3986v1 [math.RT] 22 Jan 2010

GENERALIZED VERMA MODULES AND CHARACTER FORMULAE FOR $\mathfrak{osp}(3|2m)$

BINTAO CAO AND LI LUO

Abstract. The character formula of any finite dimensional irreducible module for Lie superalgebra $\mathfrak{osp}(3|2m)$ is obtained in terms of characters of generalized Verma modules.

1. Introduction

1.1. It was already clear from the foundation papers [6, 7, 8] by Kac that finding the character formulae of the so-called atypical irreducible modules is one of the most challenging problems in the theory of Lie superalgebras.

The finite-dimensional irreducible modules of the Lie superalgebras $D(2,1;\alpha)$ and $C(n)$ were understood by van der Jeugt in [15, 16] long ago. In the case of $\mathfrak{gl}(m|n)$, the problem was first solved by Serganova in [11], where a Kazhdan-Lusztig theory was developed. Later on, Brundan [1] discovered a remarkable connection between the general linear superalgebra and the quantum group of $\mathfrak{gl}_\infty$, by which the Kazhdan-Lusztig polynomials were computed quite directly. This approach was further developed in [13, 3].

Very recently, Gruson and Serganova [5], and, Cheng, Lam and Wang [2] provided two independent and different approaches to the finite-dimensional irreducible $\mathfrak{osp}$-characters.

1.2. At the same time, the study of generalized Verma modules for ortho-symplectic Lie superalgebra $\mathfrak{osp}(n|2)$ has recently enjoyed considerable attention in [9] by Luo and in [14] by Su and Zhang. This effort is not only possibly useful to describe the finite-dimensional irreducible $\mathfrak{osp}$-characters by another approach completely different from [5, 2] but also a first step toward studying category $\mathcal{O}$.

Choose the distinguished Borel subalgebra for a basic classical Lie superalgebra and consider the maximal parabolic subalgebra obtained by removing the unique odd simple root. The generalized Verma modules are those induced from finite dimensional irreducible Verma modules over the parabolic subalgebra. In the case of types A and C, these modules coincide the Kac modules, which can be interpreted as cohomology groups of bundles over the flag supermanifold. However, in the case of types B and D, the Kac modules are not the generalized Verma modules but their maximal finite-dimensional quotients. This present paper, as well as the papers [9, 14], implies that the generalized Verma modules also play important roles in the representation theory for Lie superalgebras.

2000 *Mathematics Subject Classification.* Primary 17B10, 17B37.

Key words and phrases. ortho-symplectic superalgebra, generalized Verma module, character formula, tensor module.

1.3. Influenced by the work [9], here we shall also study any finite-dimensional irreducible module L_λ from $L_\mu \otimes \mathbb{C}^{3|2m}$ where $\mathbb{C}^{3|2m}$ is the natural representation and L_μ is another irreducible module known by induction. We point out that the module $L_\mu \otimes \mathbb{C}^{3|2m}$ may not be completely reducible. Thus we should give a more detailed argument for the blocks of modules appearing in $L_\mu \otimes \mathbb{C}^{3|2m}$.

Another key point of this work is how to express the trivial character $\mathrm{ch}\, L_0 = 1$ in terms of characters of generalized Verma modules, for which a nontrivial combinatorial identity is proved. This combinatorial identity is formulated by a certain subset of Weyl group.

1.4. The purpose of the present paper is to express the character formulae for all finite dimensional irreducible modules of $\mathfrak{osp}(3|2m)$ in terms of characters of generalized Verma modules, which are known clearly. Our main results are Theorem 3.1 and Theorem 5.2.

The paper is organized as follows. In Section 2, we present some background material on $\mathfrak{osp}(3|2m)$. In Section 3, the character formulae for tail atypical weights are obtained. Section 4 is devoted to study the information of tensor modules. The result for non-tail atypical weight is given in Section 5.

2. PRELIMINARIES

We shall simplify $\mathfrak{osp}(3|2m)$ to $\mathfrak{g}$ and work over the field $\mathbb{C}$ of complex numbers throughout the paper.

2.1. **Distinguished simple roots system.** Choose the distinguished Borel subalgebra [6] for $\mathfrak{g}$. Then the set of simple roots is

$$\Pi = \{\delta_1 - \delta_2, \delta_2 - \delta_3, \ldots, \delta_{m-1} - \delta_m, \delta_m - \epsilon, \epsilon\}. \tag{2.1}$$

The set of positive even roots and odd roots are

$$\Delta_0^+ = \{\epsilon, \delta_i \pm \delta_j, 2\delta_k \mid 1 \le i < j \le m, 1 \le k \le m\} \tag{2.2}$$

and

$$\Delta_1^+ = \{\delta_i, \delta_i \pm \epsilon \mid 1 \le i \le m\}, \tag{2.3}$$

respectively.

The distinguished Dynkin diagram is given as follows:



(Figure 1)

The bilinear form $(\cdot,\cdot)$ on $\mathfrak{h}^*$ is defined by

$$(\delta_i, \delta_j) = \left\{ \begin{array}{ll} -1, & (i = j) \\ 0, & (i \neq j) \end{array} \right. , \quad (\delta_i, \epsilon) = 0, \quad (\epsilon, \epsilon) = 1 \quad \text{for} \quad 1 \le i, j \le m. \tag{2.4}$$

2.2. **$\mathbb{Z}$-grading and parabolic subalgebras.** The Lie superalgebra $\mathfrak{g}$ admits a $\mathbb{Z}_2$-consistent $\mathbb{Z}$-grading

$$\mathfrak{g} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_2, \tag{2.5}$$

where $\mathfrak{g}_0 \cong \mathfrak{gl}(m) \oplus \mathfrak{sl}(2)$ is spanned by $\mathfrak{h}$, $\mathfrak{g}_{\pm\epsilon}$ and $\mathfrak{g}_{\pm(\delta_i-\delta_j)}$ $(1 \le i < j \le m)$; $\mathfrak{g}_{\pm 1}$ is spanned by $\mathfrak{g}_{\pm\alpha}(\alpha \in \Delta_{\bar{1}}^+)$; and $\mathfrak{g}_{\pm 2}$ is spanned by $\mathfrak{g}_{\pm(\delta_i+\delta_j)}$ $(1 \le i \le j \le m)$. There are two parabolic subalgebras

$$\mathfrak{u} = \mathfrak{g}_1 \oplus \mathfrak{g}_2 \quad \text{with} \quad \mathfrak{u}_{\bar{0}} = \mathfrak{g}_2 \text{ and } \mathfrak{u}_{\bar{1}} = \mathfrak{g}_1 \tag{2.6}$$

and

$$\mathfrak{u}^- = \mathfrak{g}_{-1} \oplus \mathfrak{g}_{-2} \quad \text{with} \quad \mathfrak{u}^-_{\bar{0}} = \mathfrak{g}_- \text{ and } \mathfrak{u}^-_{\bar{1}} = \mathfrak{g}_{-1}. \tag{2.7}$$

2.3. **Dominant integral weights.** An element in $\mathfrak{h}^*$ is called a *weight.* A weight $\lambda \in \mathfrak{h}^*$ will be written in terms of the $\delta\epsilon$-basis as

$$\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_m; \lambda_0) = \sum_{i=1}^{m} \lambda_i \delta_i + \lambda_0 \epsilon. \tag{2.8}$$

Define

$$\text{ht } \lambda = \sum_{i=0}^{m} |\lambda_i|, \tag{2.9}$$

and call it the height of λ.

A weight λ is called *integral* if $\lambda_0, \lambda_1, \ldots, \lambda_m \in \mathbb{Z}$.

For any weight $\lambda \in \mathfrak{h}^*$, there exists an irreducible module L_λ with highest weight λ.

Theorem 2.1. (Kac [6]**)** *For any integral weight λ, the irreducible module L_λ is finite dimensional if and only if*

$$\begin{cases} \lambda_0, \lambda_1, \ldots, \lambda_m \in \mathbb{Z}_{\ge 0}, \\ \lambda_1 \ge \lambda_2 \ge \cdots \ge \lambda_{m-1} \ge \lambda_m, \\ \lambda_0 = 0 \quad \textit{if} \quad \lambda_m = 0. \end{cases} \tag{2.10}$$

Denote

$$\mathcal{P} = \{\lambda \in \mathfrak{h}^* \mid \lambda \text{ is integral and satisfies the conditions in Theorem 2.1}\}. \tag{2.11}$$

The weights in $\mathcal{P}$ are called *dominant integral.*

2.4. **Generalized Verma modules.** For any $\lambda \in \mathfrak{h}^*$, we denote by $L_\lambda^{(0)}$ the irreducible $\mathfrak{g}_0$- module with highest weight λ. Denote

$$\mathcal{P}^{(0)} = \{\lambda \in \mathfrak{h}^* \mid L_\lambda^{(0)} \text{ is finite-dimensional}\}. \tag{2.12}$$

It is obvious that $\mathcal{P} \subset \mathcal{P}^{(0)}$.

For any $\lambda \in \mathcal{P}^{(0)}$, extend $L_\lambda^{(0)}$ to a $\mathfrak{g}_0 \oplus \mathfrak{u}$-module by putting $\mathfrak{u}L_\lambda^{(0)} = 0$. Then the *generalized Verma module* M_λ is defined as the induced module

$$M_\lambda = \text{Ind}_{\mathfrak{g}_0 \oplus \mathfrak{u}}^{\mathfrak{g}} L_\lambda^{(0)} \cong U(\mathfrak{u}^-) \otimes_{\mathbb{C}} L_\lambda^{(0)}. \tag{2.13}$$

It is clear that L_λ is the unique irreducible quotient module of M_λ.

2.5. **Atypical weights.** Let ρ_0 (resp. ρ_1) be half the sum of positive even (resp. odd) roots, and let $\rho = \rho_0 - \rho_1$. Then

$$\rho = (m - \frac{3}{2})\delta_1 + (m - \frac{5}{2})\delta_2 + \cdots + \frac{1}{2}\delta_{m-1} - \frac{1}{2}\delta_m + \frac{1}{2}\epsilon. \tag{2.14}$$

A weight $\lambda \in \mathfrak{h}^*$ is called *atypical* if there is a positive odd root $\alpha \in \bar{\Delta}_1^+$ such that

$$(\lambda + \rho, \alpha) = 0. \tag{2.15}$$

Sometimes we also call it α*-atypical* to emphasize the odd root α. Otherwise, we call λ *typical*.

Note that if $\lambda \in \mathcal{P}$ is $(\delta_t - \epsilon)$-atypical for some $1 \leq t \leq m$, then it must be that $t = m$ and $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_{m-1} \geq \lambda_m = \lambda_0 = 0$. Such a weight λ is called a *tail atypical weight.* A weight $\lambda \in \mathcal{P}$ which is not a tail atypical weight, is called a *non-tail atypical weight.*

For convenience, sometimes we use the ρ*-translated* notation $\widetilde{\lambda}$ of a weight λ, which is defined by

$$\widetilde{\lambda} = \lambda + \rho = (\widetilde{\lambda}_1, \widetilde{\lambda}_2, \ldots, \widetilde{\lambda}_m; \widetilde{\lambda}_0) \tag{2.16}$$

Assume $\lambda \in \mathfrak{h}^*$ is α-atypical with $\alpha = \delta_t + \epsilon$ or $\delta_t - \epsilon$. Denote

$$\overline{\lambda} = [|\widetilde{\lambda}_1|, \ldots, |\widetilde{\lambda}_{t-1}|, |\widetilde{\lambda}_{t+1}|, \ldots, |\widetilde{\lambda}_m|], \tag{2.17}$$

which is called the *atypical type* of λ.

2.6. **Weyl group.** Let $\mathcal{W}$ be the Weyl group of $\mathfrak{g}$ (i.e. the Weyl group of Lie algebra $\mathfrak{g}_{\bar{0}}$) and $\ell : \mathcal{W} \to \mathbb{N}$ be the length function. Let $S_k \ltimes \mathbb{Z}_2^k \subset \mathcal{W}$ be the Weyl group of type C_k $(1 \leq k \leq m)$. That is, elements in S_k permute the coefficients of $\delta_1, \ldots, \delta_k$ and elements in $\mathbb{Z}_2^k$ show how to change the signs of the coefficients of $\delta_1, \ldots, \delta_k$.

For any $1 \leq k \leq m$, denote

$$\Gamma_k := \{\sigma \in S_k \ltimes \mathbb{Z}_2^k \mid \sigma(\mu)_1 > \sigma(\mu)_2 > \cdots > \sigma(\mu)_k\}, \tag{2.18}$$

where $\mu = k\delta_1 + (k-1)\delta_2 + \cdots + \delta_k$ and $\sigma(\mu)_i$ is the coefficient of δ_i in $\sigma(\mu)$.

Remark 2.2. Note that $\sigma = \omega \cdot \varepsilon \in \Gamma_{m-1}$ is determined by ε uniquely, where $\omega \in S_{m-1}$ and $\varepsilon \in \mathbb{Z}_2^{m-1}$. Hence there is a 1-1 correspondence between the elements in Γ_{m-1} and the self-conjugate partitions with length smaller than m via

$$\mu = (j_p - 1, \ldots, j_1 - 1 | j_p - 1, \ldots, j_1 - 1) \longmapsto \sigma_\mu = \omega_\mu \cdot \varepsilon_\mu, \tag{2.19}$$

where $1 \leq j_1 \leq \cdots \leq m-1$ and $\varepsilon_\mu = (\varepsilon_{\mu,1}, \ldots, \varepsilon_{\mu,m-1})$ with

$$\varepsilon_{\mu,s} = \begin{cases} -1 & \text{if } \ s \in \{m - j_1, \ldots, m - j_p\} \\ 1 & \text{otherwise.} \end{cases} \tag{2.20}$$

2.7. **Central characters and blocks.** Denote by $Z(\mathfrak{g})$ the central of the universal enveloping algebra $U(\mathfrak{g})$. Recall that a *central character* defined by $\lambda \in \mathfrak{h}^*$ is a homomorphism $\chi_\lambda : Z(\mathfrak{g}) \to \mathbb{C}$ such that each element $z \in Z(\mathfrak{g})$ acts on L_λ as the scaler $\chi_\lambda(z)$.

Lemma 2.3. (See. [12]) *For any $\lambda, \mu \in \mathcal{P}^{(0)}$, $\chi_\lambda = \chi_\mu$ implies that $\overline{\lambda} = \overline{\mu}$ if λ, μ are both atypical or that there exists $w \in \mathcal{W}$ such that $\widetilde{\lambda} = w(\widetilde{\mu})$ if λ, μ are both typical.*

Suppose $\lambda \in \mathcal{P}$ is typical, then for any $\mu \in \mathcal{P}$, $\chi_\lambda = \chi_\mu$ if and only if $\lambda = \mu$.

For any atypical weight $\lambda \in \mathcal{P}$, we denote by λ^T the unique tail atypical weight with $\chi_\lambda = \chi_{\lambda^T}$. It is obvious that $\lambda = \lambda^T$ if λ is a tail atypical weight. For a non-tail atypical weight $\lambda \in \mathcal{P}$, one can check that

$$\lambda^T = \sum_{i=1}^{k-1} \lambda_i \delta_i + \sum_{i=k}^{m-1} \lambda_{i+1} \delta_i \quad \text{if } \lambda \text{ is } (\delta_k + \epsilon)\text{-atypical.} \tag{2.21}$$

For any tail atypical weight λ, denote

$$B_\lambda = \{\mu \in \mathcal{P} \mid \mu^T = \lambda\}. \tag{2.22}$$

Clearly,

$$\mathcal{P} = \bigcup_{\lambda \text{ tail atypical}} B_\lambda, \quad \text{which is a disjoint union.} \tag{2.23}$$

Take any non-tail $(\delta_k + \epsilon)$-atypical weight λ $(1 \leq k \leq m)$ and assume that t is the maximal number with $\lambda_t = \lambda_k$. Define

$$\varphi(\lambda) = \begin{cases} \lambda - \delta_m = \lambda^T, & \text{if } \lambda_0 = 0 \\ \lambda - \sum_{i=k}^{t} \delta_i - (t-k+1)\epsilon, & \text{otherwise.} \end{cases} \tag{2.24}$$

It is obvious that $\varphi : \mathcal{P} \setminus \{\text{tail atypical weights}\} \to \mathcal{P}$ is injective. Moreover, one can check that $\varphi(\mu) \in B_\lambda$ for any tail typical weight λ and any $\mu \in B_\lambda$.

2.8. **Characters for typical weights.**

Theorem 2.4. (Kac[7]) A weight $\lambda \in \mathcal{P}$ is typical if and only if

$$ch\ L_\lambda = \frac{\prod_{\alpha \in \Delta_1^+} (e^{\frac{\alpha}{2}} + e^{-\frac{\alpha}{2}})}{\prod_{\alpha \in \Delta_0^+} (e^{\frac{\alpha}{2}} - e^{-\frac{\alpha}{2}})} \sum_{w \in \mathcal{W}} (-1)^{l(w)} e^{w(\lambda+\rho)}. \tag{2.25}$$

We rewrite this theorem as the following lemma.

Lemma 2.5. *A weight $\lambda \in \mathcal{P}$ is typical if and only if*

$$ch\ L_\lambda = \sum_{\sigma \in \Gamma_m} (-1)^{\ell(\sigma)} ch\ M_{\sigma(\lambda+\rho)-\rho}. \tag{2.26}$$

Proof. By Theorem 2.4, one has that $\lambda \in \mathcal{P}$ is typical if and only if

$$ch\ L_\lambda = \frac{\prod\limits_{\alpha \in \Delta_1^+} (1 + e^{-\alpha})}{\prod\limits_{1 \leq i \leq j \leq m} (1 - e^{-(\delta_i + \delta_j)})} \frac{\sum\limits_{w \in \mathcal{W}} (-1)^{l(w)} e^{w(\lambda+\rho)-\rho}}{(1 - e^{-\epsilon}) \prod\limits_{1 \leq i < j \leq m} (1 - e^{-(\delta_i - \delta_j)})}. \tag{2.27}$$

For convenience, we denote $\mu' = \sum_{j=1}^m \mu_j \delta_j$ with $\mu = \sum_{j=1}^m \mu_j \delta_j + \mu_0 \epsilon$, and denote $\rho_{\mathfrak{gl}(m)} = \sum_{j=1}^m (m-j)\delta_j$ being half the sum of positive roots of $\mathfrak{sl}(m)$. Since $\mathcal{W} = \mathcal{W}_{C_m} \times \mathcal{W}_{B_1}$, we have

$$\sum_{w \in \mathcal{W}} (-1)^{l(w)} e^{w(\lambda+\rho)-\rho} = (e^{\lambda_0 \epsilon} - e^{-(\lambda_0+1)\epsilon}) \sum_{w \in S_m \ltimes \mathbb{Z}_2^m} (-1)^{l(w)} e^{w(\lambda'+\rho')-\rho'}. \tag{2.28}$$

Thus

$$\begin{aligned}
ch\ L_\lambda &= \frac{\prod\limits_{\alpha\in\Delta_1^+}(1+e^{-\alpha})}{\prod\limits_{1\le i\le j\le m}(1-e^{-(\delta_i+\delta_j)})}\sum_{j=0}^{2\lambda_0} e^{(\lambda_0-j)\epsilon}\frac{\sum\limits_{w\in S_m\ltimes\mathbb{Z}_2^m}(-1)^{l(w)}e^{w(\lambda'+\rho')-\rho'}}{\prod\limits_{1\le i<j\le m}(1-e^{-(\delta_i-\delta_j)})}\\
&= \frac{\prod\limits_{\alpha\in\Delta_1^+}(1+e^{-\alpha})}{\prod\limits_{1\le i\le j\le m}(1-e^{-(\delta_i+\delta_j)})}\sum_{j=0}^{2\lambda_0} e^{(\lambda_0-j)\epsilon}\sum_{\sigma\in\Gamma_m}(-1)^{l(\sigma)}\frac{\sum\limits_{w\in S_m}(-1)^{l(w)}e^{w(\sigma(\lambda'+\rho')-\rho'+\rho_{\mathfrak{gl}(m)})}}{e^{\rho_{\mathfrak{gl}(m)}}\prod\limits_{1\le i<j\le m}(1-e^{-(\delta_i-\delta_j)})}\\
&= \frac{\prod\limits_{\alpha\in\Delta_1^+}(1+e^{-\alpha})}{\prod\limits_{1\le i\le j\le m}(1-e^{-(\delta_i+\delta_j)})}\sum_{\sigma\in\Gamma_m}(-1)^{l(\sigma)}ch\ L^{(0)}_{\sigma(\lambda+\rho)-\rho}\\
&= \sum_{\sigma\in\Gamma_m}(-1)^{l(\sigma)}ch\ M_{\sigma(\lambda+\rho)-\rho}
\end{aligned}$$

□

2.9. **Cohomology and character.** The space of q-dimensional cochains of the Lie superalgebra $\mathfrak{u}=\mathfrak{u}_{\bar{0}}\otimes\mathfrak{u}_{\bar{1}}$ with coefficients in the module L_λ is given by

$$C^q(\mathfrak{u};L_\lambda)=\bigoplus_{q_0+q_1=q}\mathrm{Hom}(\wedge^{q_0}\mathfrak{u}_{\bar{0}}\otimes S^{q_1}\mathfrak{u}_{\bar{1}},L_\lambda). \tag{2.29}$$

The differential $d:C^q(\mathfrak{u};L_\lambda)\to C^{q+1}(\mathfrak{u};L_\lambda)$ is defined by

$$\begin{gathered}
dc(\xi_1,\dots,\xi_{q_0},\eta_1,\dots,\eta_{q_1})\\
=\textstyle\sum_{1\le s\le t\le q_0}(-1)^{s+t-1}c([\xi_s,\xi_t],\xi_1,\dots,\widehat{\xi_s},\dots,\widehat{\xi_t},\dots,\xi_{q_0},\eta_1,\dots,\eta_{q_1})\\
+\textstyle\sum_{s=1}^{q_0}\sum_{t=1}^{q_1}(-1)^{s-1}c(\xi_1,\dots,\widehat{\xi_s},\dots,\xi_{q_0},[\xi_s,\eta_t],\eta_1,\dots,\widehat{\eta_t},\dots,\eta_{q_1})\\
+\textstyle\sum_{1\le s\le t\le q_1}c([\eta_s,\eta_t],\xi_1,\dots,\xi_{q_0},\eta_1,\dots,\widehat{\eta_s},\dots,\widehat{\eta_t},\dots,\eta_{q_1})\\
+\textstyle\sum_{s=1}^{q_0}(-1)^s\xi_sc(\xi_1,\dots,\widehat{\xi_s},\dots,\xi_{q_0},\eta_1,\dots,\eta_{q_1})\\
+(-1)^{q_0-1}\textstyle\sum_{s=1}^{q_1}\eta_sc(\xi_1,\dots,\xi_{q_0},\eta_1,\dots,\widehat{\eta_s},\dots,\eta_{q_1})
\end{gathered}\tag{2.30}$$

where $c\in C^q(\mathfrak{u};L_\lambda)$, $\xi_1,\dots,\xi_{q_0}\in\mathfrak{u}_{\bar{0}}$, $\eta_1,\dots,\eta_{q_1}\in\mathfrak{u}_{\bar{1}}$.

The cohomology of $\mathfrak{u}$ with coefficients in the module L_λ is the cohomology group of the complex $C=(\{C^q(\mathfrak{u};L_\lambda)\},d)$, and it is denoted by $H^q(\mathfrak{u},L_\lambda)$.

Imitating the argument in [9], one can obtain

$$\mathrm{ch}\ L_\lambda=\mathrm{ch}\ M_\lambda+\sum_{\nu\prec\lambda}\sum_{i=0}^{\infty}(-1)^i[H^i(\mathfrak{u},L_\lambda):L_\nu^{(0)}]\mathrm{ch}\ M_\nu, \tag{2.31}$$

where $\nu\prec\lambda$ means that $\chi_\lambda=\chi_\nu$ and $\nu<\lambda$ (i.e. $\lambda-\mu$ is a $\mathbb{Z}_{\ge0}$-linear sum of positive roots), and $[H^i(\mathfrak{u},L_\lambda):L_\nu^{(0)}]$ is the multiplicity of $L_\nu^{(0)}$ in the cohomology group regarded as a $\mathfrak{g}_0$-module.

Remark 2.6. In the above expression, any coefficient of ch M_ν is an integral number. In particular, the coefficient of ch M_λ is 1.

3. Character formulae for tail atypical weights

3.1. **Natural module.** Consider the natural $\mathfrak{g}$-module $L_{\delta_1} \simeq \mathbb{C}^{3|2m}$. It is known that ${L_{\delta_1}}^* \simeq L_{\delta_1}$ and that the set of weights of L_{δ_1} is $\{0, \pm\delta_1, \dots, \pm\delta_m, \pm\epsilon\}$. Moreover, as a $\mathfrak{g}_0$-module,

$$L_{\delta_1} \simeq L^{(0)}_{\delta_1} \oplus L^{(0)}_{-\delta_m} \oplus L^{(0)}_{\epsilon}. \tag{3.1}$$

3.2. **Weights set $\mathcal{P}_\lambda$ and $\mathcal{P}^{(0)}_\lambda$.** For any $\lambda \in \mathcal{P}$, denote

$$\mathcal{P}_\lambda = \mathcal{P}_{\lambda^+} \cup \mathcal{P}_{\lambda^-} \tag{3.2}$$

where

$$\mathcal{P}_{\lambda^+} = \begin{cases} \{\lambda+\delta_1, \lambda+\delta_2, \dots, \lambda+\delta_m, \lambda+\epsilon\} \cap \mathcal{P}, & (\lambda_0 = 0); \\ \{\lambda+\delta_1, \lambda+\delta_2, \dots, \lambda+\delta_m, \lambda, \lambda+\epsilon\} \cap \mathcal{P}, & (\lambda_0 \neq 0). \end{cases} \tag{3.3}$$

and

$$\mathcal{P}_{\lambda^-} = \begin{cases} \{\lambda-\delta_1, \lambda-\delta_2, \dots, \lambda-\delta_m, \lambda-\epsilon\} \cap \mathcal{P}, & (\lambda_0 = 0); \\ \{\lambda-\delta_1, \lambda-\delta_2, \dots, \lambda-\delta_m, \lambda, \lambda-\epsilon\} \cap \mathcal{P}, & (\lambda_0 \neq 0). \end{cases} \tag{3.4}$$

Let $\mathcal{P}^{(0)}$ take the place of $\mathcal{P}$ above, then there come the definitions of $P^{(0)}_\lambda$, $\mathcal{P}^{(0)}_{\lambda^+}$ and $\mathcal{P}^{(0)}_{\lambda^+}$.

The following statement is standard in the theory of the classical finite-dimensional semisimple Lie algebras: for any $\lambda \in \mathcal{P}^{(0)}$,

$$L^{(0)}_\lambda \otimes (L^{(0)}_{\delta_1} \oplus L^{(0)}_{-\delta_m} \oplus L^{(0)}_{\epsilon}) = \bigoplus_{\mu \in P^{(0)}_\lambda} L^{(0)}_\mu. \tag{3.5}$$

3.3. **Character of $M_\lambda \otimes L_{\delta_1}$.** Thanks to (3.1) and (3.5), we have that for any $\lambda \in \mathcal{P}^{(0)}$, as a $\mathfrak{g}_0$-module,

$$\begin{aligned} M_\lambda \otimes L_\delta &= (U(\mathfrak{u}^-) \otimes_{\mathbb{C}} L^{(0)}_\lambda) \otimes (L^{(0)}_{\delta_1} \oplus L^{(0)}_{-\delta_m} \oplus L^{(0)}_{\epsilon}) \\ &\cong U(\mathfrak{u}^-) \otimes_{\mathbb{C}} (L^{(0)}_\lambda \otimes (L^{(0)}_{\delta_1} \oplus L^{(0)}_{-\delta_m} \oplus L^{(0)}_{\epsilon})) \\ &= U(\mathfrak{u}^-) \otimes_{\mathbb{C}} (\bigoplus_{\mu \in \mathcal{P}^{(0)}_\lambda} L^{(0)}_\mu) \\ &= \bigoplus_{\mu \in \mathcal{P}^{(0)}_\lambda} (U(\mathfrak{u}^-) \otimes_{\mathbb{C}} L^{(0)}_\mu) \\ &= \bigoplus_{\mu \in \mathcal{P}^{(0)}_\lambda} M_\mu. \end{aligned} \tag{3.6}$$

Hence

$$\text{ch } M_\lambda \otimes L_\delta = \sum_{\mu \in \mathcal{P}^{(0)}_\lambda} \text{ch } M_\mu. \tag{3.7}$$

3.4. **Convention.** From here on, we will always simplify the character ch V to V. It cannot confuse us by context.

3.5. **Character formulae.** Firstly, we will recall some notations and facts about symmetric functions. One can find more material in [10].

Denote $A_\lambda = \det(x_j^{\lambda_i})_{1\le i,j\le m}$ for $\lambda = (\lambda_1, \dots, \lambda_m) \in \mathbb{Z}^m$ and $S_\lambda = A_{\lambda+\rho_{\mathfrak{gl}(m)}}/A_{\rho_{\mathfrak{gl}(m)}}$, where $\rho_{\mathfrak{gl}(m)} = \sum_{j=1}^m (m-j)\delta_j$. Note that λ is not necessary a partition, and if so then we will get back to the traditional definition of Schur function S_λ. Thus $S_\lambda \neq 0$ if and only if there exists $w \in S_m$ satisfying $\lambda_{\sigma(1)} - \sigma(1) + 1 \geq \cdots \geq \lambda_{\sigma(m)} - \sigma(m) + m$. Equivalently, $S_\lambda = 0$ if and only if there exist $i, j \in \{1, \dots, m\}$ satisfying that $\lambda_i - i = \lambda_j - j$.

A direct calculation shows that $S_\lambda = \det(h_{\lambda_i+l-i+j})/e_m^l$, where $l \in \mathbb{Z}_{\geq 0}$ is any nonnegative integer such that $\lambda_i + l - i + m \geq 0$ for all $i \in \{1, \dots, m\}$. Here $h_l = h_l(x_1, \dots, x_m)$ is the l-th complete symmetric polynomials (i.e. the sum of all monomials of total degree l in the variables $x_1, \dots, x_m$), and $e_l = e_l(x_1, \dots, x_m)$ is the l-th elementary symmetric polynomials in the variables $x_1, \dots, x_m$. We set $h_l = e_l = 0$ if $l < 0$ for convenience.

For each $\sigma \in \Gamma_{m-1}$ (cf. (2.18)), define

(3.8)
$$\flat(\sigma, \lambda) = \begin{cases} i, & \text{if } i \in \{1, 2, \dots, m-1\} \text{ is the minimal number such that } \sigma(\lambda+\rho)_i < -1; \\ m, & \text{if } \sigma(\lambda+\rho)_{m-1} > -1. \end{cases}$$

Write

$$\tau_i := (i, i+1, i+2, \dots, m) \in S_m. \tag{3.9}$$

Theorem 3.1. *i). If λ is a tail atypical dominant integral weight, then*

$$L_\lambda = \sum_{\sigma\in\Gamma_{m-1}} \sum_{i=\flat(\sigma,\lambda)}^{m} \sum_{j=\max\{0, \frac{1}{2}-\sigma(\lambda+\rho)_{i-1}\}}^{-\frac{3}{2}-\tau_i\sigma(\lambda+\rho)_{i+1}} (-1)^{\ell(\tau_i\sigma)+j} M_{\tau_i\sigma(\lambda+\rho)-\rho+j\epsilon-j\delta_i}, \tag{3.10}$$

where $\sigma(\lambda+\rho)_0 = \infty$ and $\tau_m\sigma(\lambda+\rho)_{m+1} = -\infty$ for convenience.

ii). If $\lambda = \sum_{j=1}^m \lambda_j\delta_j + \lambda_0\epsilon$ is a $(\delta_k + \epsilon)$-atypical dominant integral weight with $\lambda_k = \cdots \lambda_m = 1$ and $\lambda_0 = m - k$ (i.e. λ is a non-tail atypical weight with $\varphi(\lambda) = \lambda^T$), then

$$L_\lambda = L_{\lambda^T} + \sum_{\sigma\in\Gamma_m} (-1)^{\ell(\sigma)} ch\ M_{\sigma(\lambda+\rho)-\rho}. \tag{3.11}$$

where L_{λ^T} has been got by i).

We will give several lemmas first.

Denote F_λ the right side of (3.10) or (3.11).

If λ is a $(\delta_m - \epsilon)$-atypical integral dominant weight of $\mathfrak{g}$, we define

(3.12)
$$C_\lambda = \frac{\prod_{i=1}^m (1+x_iy) \prod_{i=1}^m (1+x_iy^{-1})}{\prod_{i=1}^m (1-x_i) \prod_{1\le i<j\le m} (1-x_ix_j)} \sum_{j=0}^{\infty} \sum_{\mu} (-1)^{j+T(\mu)} S_{(j,\mu)} \sum_{s=0}^{2j} y^{j-s},$$

where μ runs over all partitions in forms $(j_p - 1, \dots, j_1 - 1 | j_p - 1, \dots, j_1 - 1) + (\lambda_{m-j_p}, \dots, \lambda_{m-j_1}, -\lambda_{m-r_1}, \dots, -\lambda_{m-r_{m-1-p}})$ for any nonnegative integer $p < m$ and strictly increasing p-tuples $(j_1, \dots, j_p) \in \{1, \dots, m-1\}^p$. Here the entries of the strictly increasing $(m-1-p)$-tuples $(r_1, \dots, r_{m-1-p})$ are in the set $\{1, \dots, m-1\}\backslash\{j_1, \dots, j_p\}$, and $T(\mu) = j_1 + \cdots + j_p$.

If λ is a $(\delta_m+\epsilon)$-atypical dominant integral weight with $\lambda_m=1$ and $\lambda_0=0$, we define

(3.13)
$$C_\lambda = C_{\lambda-\delta_m} + \frac{\prod_{i=1}^{m}(1+x_iy)\prod_{i=1}^{m}(1+x_iy^{-1})}{\prod_{i=1}^{m}(1-x_i)\prod_{1\le i<j\le m}(1-x_ix_j)}\sum_{\mu}(-1)^{T(\mu)}(S_{(-1,\mu)}-S_{(0,\mu)}),$$

where the definitions of μ and $T(\mu)$ are as the same as in (3.12).

Lemma 3.2. *$C_\lambda = F_\lambda$ if one sets $x_i = e^{-\delta_i}$ and $y=e^{\epsilon}$.*

Proof. For convenience, we denote $\lambda^c = \lambda_m\delta_1+\cdots+\lambda_1\delta_m$ and $\rho^c = -\frac{1}{2}\delta_1+\frac{1}{2}\delta_2+\cdots+(m-\frac{3}{2})\delta_m$. Then

$$\Gamma_{m-1} = \{\sigma\in S_{m-1}\ltimes\mathbb{Z}_2^{m-1}\mid \sigma(\lambda^c+\rho^c)_2>\cdots>\sigma(\lambda^c+\rho^c)_m\}. \tag{3.14}$$

Take

$$\upsilon = (j_p-1,\ldots,j_1-1|j_p-1,\ldots,j_1-1) \tag{3.15}$$

(3.16)
$$\mu = \upsilon + (\lambda_{m-j_p},\ldots,\lambda_{m-j_1},-\lambda_{m-r_1},\ldots,-\lambda_{m-r_{m-1-p}}),$$

and

$$\sigma_\mu = \omega_\upsilon\cdot\varepsilon_\upsilon \tag{3.17}$$

for $\omega_\upsilon\in S_{m-1}$, $\varepsilon_\upsilon\in\mathbb{Z}_2^{m-1}$ and

$$\varepsilon_{\upsilon,s} = \begin{cases} -1 & \text{if } s\in\{m-j_1,\ldots,m-j_p\} \\ 1 & \text{if } s\in\{m-r_1,\ldots,m-r_{m-1-p}\}\end{cases}. \tag{3.18}$$

One easily shows that

$$(-1)^{j+T(\mu)}S_{(j,\mu)} = (-1)^{j+l(\sigma_\mu)}S_{\rho^c-\sigma_\mu(\lambda^c+\rho^c)+j\delta_1}. \tag{3.19}$$

Since there is a 1-1 correspondence between the elements in Γ_{m-1} and the strict increasing $p-$tuples $(j_1,\ldots,j_p)\in\{1,\ldots,m-1\}^p$ (see Remak 2.2), where p runs over the set $\{1,\ldots,m-1\}$, we get that

$$\sum_\mu(-1)^{j+T(\mu)}S_{(j,\mu)} = \sum_{\sigma\in\Gamma_{m-1}}(-1)^{j+l(\sigma)}S_{\rho^c-\sigma(\lambda^c+\rho^c)+j\delta_1}. \tag{3.20}$$

For an $m-$tuple $(\nu_1,\ldots,\nu_m)$, one has

$$S_{(\nu_1,\ldots,\nu_m)}(x_1,\ldots,x_m) = S_{(-\nu_m,\ldots,-\nu_1)}(x_1^{-1},\ldots,x_m^{-1}). \tag{3.21}$$

Thus

$$S_{\rho^c-\sigma(\lambda^c+\rho^c)+j\delta_1}(x_1,\ldots,x_m) = S_{\sigma(\lambda+\rho)-\rho-j\delta_m}(x_1^{-1},\ldots,x_m^{-m}). \tag{3.22}$$

Furthermore, it is well known that $S_\eta = (-1)^{l(w)}S_\theta$ if $\eta+\rho_{\mathfrak{gl}(m)} = w(\theta+\rho_{\mathfrak{gl}(m)})$, where $w\in S_m$. One checks that $S_{\sigma(\lambda+\rho)-\rho-j\delta_m}=0$ if $j\notin\{\max\{0,\frac{1}{2}-\sigma(\lambda+\rho)_{i-1}\},\ldots,-\frac{1}{2}-\tau_i\sigma(\lambda+\rho)_{i+1}\}$ for $i\in\{\flat(\sigma,\lambda),\ldots,m\}$ in (3.10). These complete the proof. ☐

Lemma 3.3. *The following identity holds:*

$$\frac{\prod_{i=1}^{m}(1-x_i)\prod_{1\le i<j\le m}(1-x_ix_j)}{\prod_{i=1}^{m}(1+x_iy)\prod_{i=1}^{m}(1+x_iy^{-1})}=\sum_{j=0}^{\infty}\sum_{\mu}(-1)^{j+\frac{|\mu|+p(\mu)}{2}}S_{(j,\mu)}\sum_{s=0}^{2j}y^{j-s}, \tag{3.23}$$

where μ *runs over all self-conjugate partitions* $\mu=(\alpha_1,\ldots,\alpha_p|\alpha_1,\ldots,\alpha_p)$ *with* $\alpha_1\le m-2$. *Here* $p(\mu)=p$ *and* $S_{(j,\mu)}=S_{(j,\mu)}(x_1,\ldots,x_m)$.

Proof. One shows that

$$\prod_{i=1}^{m}(1+x_iy)\prod_{i=1}^{m}(1+x_iy^{-1})=\sum_{k=0}^{m}(\sum_{j=0}^{k}e_je_{m-k+j})y^{m-k}+\sum_{k=1}^{m}(\sum_{j=0}^{m-k}e_{k+j}e_j)y^{-k}, \tag{3.24}$$

where $e_j=e_j(x_1,\ldots,x_m)$.

We denote

$$A_j=\sum_{\substack{\mu=(\alpha_1,\ldots,\alpha_p|\alpha_1,\ldots,\alpha_p)\\ 0\le\alpha_1\le m-2}}(-1)^{\frac{|\mu|+p(\mu)}{2}}S_{(j,\mu)} \tag{3.25}$$

On the other hand, we can write $A_j=\sum_{s=0}^{m-1}a_sh_{j+s}$ by the definition of S_λ, where a_s are certain polynomials in variables $x_1,\ldots,x_m$, and $h_j=h_j(x_1,\ldots,x_m)$. We write

$$B_k=\begin{cases}\sum_{j=k}^{\infty}(-1)^jA_j & if\quad k\ge 0,\\ \sum_{j=-k}^{\infty}(-1)^jA_j & if\quad k<0.\end{cases} \tag{3.26}$$

Thus $B_k=B_{-k}$. Moreover

$$\sum_{j=0}^{\infty}\sum_{\mu}(-1)^j(-1)^{\frac{|\mu|+p(\mu)}{2}}S_{(j,\mu)}\sum_{s=0}^{2j}y^{j-s}=\sum_{k=-\infty}^{\infty}B_ky^k. \tag{3.27}$$

Hence

$$\begin{aligned}&\prod_{i=1}^{m}(1+x_iy)\prod_{i=1}^{m}(1+x_iy^{-1})(\sum_{k=-\infty}^{\infty}B_ky^k)\\ =&\sum_{l=-\infty}^{\infty}(\sum_{k=0}^{m}\sum_{j=0}^{k}e_je_{m-k+j}B_{l-m+k}+\sum_{k=1}^{m}\sum_{j=0}^{m-k}e_{k+j}e_jB_{l+k})y^l.\end{aligned} \tag{3.28}$$

Now we consider the coefficients of y^l. Since the coefficient of y^l is the same as of y^{-l}, we only need to check the case $l\ge 0$. In fact, we want to show these coefficients are 0 if $l>0$, and is $\prod_{i=1}^{m}(1-x_i)\prod_{1\le i<j\le m}^{m}(1-x_ix_j)$ if $l=0$.

Note that

(3.29)

$$\sum_{k=0}^{m}\sum_{j=0}^{k}e_je_{m-k+j}B_{l-m+k}+\sum_{k=1}^{m}\sum_{j=0}^{m-k}e_{k+j}e_jB_{l+k}=\sum_{j=0}^{m}e_j\sum_{k=0}^{m}e_{m-k}B_{l-(m-j)+k},$$

$$B_p=\sum_{j=p}^{\infty}(-1)^jA_j=\sum_{j=p}^{\infty}\sum_{s=0}^{m-1}(-1)^ja_sh_{j+s} \tag{3.30}$$

for $p \geq 0$, and

$$h_k - e_1 h_{k-1} + e_2 h_{k-2} + \cdots + (-1)^m e_m h_{h-m} = 0 \tag{3.31}$$

for $k \neq 0$. Thus the coefficient of y^l is 0 if $l \geq m$.

Let $l = m-1$, It is sufficiently to show that

$$e_m B_{-1} + e_{m-1} B_0 + \cdots + e_1 B_{m-2} + B_{m-1} = 0. \tag{3.32}$$

Since

$$e_m B_0 + e_{m-1} B_1 + \cdots + e_1 B_{m-1} + B_m = 0 \tag{3.33}$$

by(3.29)-(3.31), then (3.32) is equal to

$$A_0 = A_{-1}. \tag{3.34}$$

By induction, one shows that the equality(3.23) is equal to

$$\begin{cases} A_j = A_{-(j+1)} & \text{for } j = 0, 1, \ldots, m-2, \\ (-1)^m A_{m-1} + a_0 = \prod_{i=1}^{m} (1 - x_i) \prod_{1 \leq i < j \leq m} (1 - x_i x_j). & \end{cases} \tag{3.35}$$

Note that[10]

$$\prod_{i=1}^{m} (1 - x_i) \prod_{1 \leq i < j \leq m} (1 - x_i x_j) = \sum_{\mu} (-1)^{\frac{|\mu| + p(\mu)}{2}} S_\mu, \tag{3.36}$$

where $\mu = (\alpha_1, \ldots, \alpha_p | \alpha_1, \ldots, \alpha_p)$ and $\alpha_1 \leq m-1$. But

$$a_0 = \sum_{\mu} (-1)^{\frac{|\mu| + p(\mu)}{2}} S_\mu \tag{3.37}$$

where $\mu = (\alpha_1, \ldots, \alpha_p | \alpha_1, \ldots, \alpha_p)$ and $\alpha_1 \leq m-2$. Then

$$\begin{aligned} & \prod_{i=1}^{m} (1 - x_i) \prod_{1 \leq i < j \leq m} (1 - x_i x_j) - a_0 \\ = \; & e_m \sum_{\mu = (\alpha_1, \ldots, \alpha_p | \alpha_1, \ldots, \alpha_p)} (-1)^{\frac{|\mu| + p(\mu) + 2m}{2}} S_{(m-1, \mu)} \\ = \; & (-1)^m e_m A_{m-1}. \end{aligned} \tag{3.38}$$

Now the statement $(3.35)_2$ is hold.

Observe that

$$\begin{aligned} & (\alpha_1, \ldots, \alpha_p | \alpha_1, \ldots, \alpha_p) \\ = \; & (\alpha_1 + 1, \ldots, \alpha_p + p, \overbrace{p, \ldots, p}^{\alpha_p}, \overbrace{p-1, \ldots, p-1}^{\alpha_{p-1} - \alpha_p - 1}, \ldots, \overbrace{1, \ldots, 1}^{\alpha_1 - \alpha_2 - 1}, \overbrace{0, \ldots, 0}^{m - \alpha_1 - 2}) \end{aligned} \tag{3.39}$$

Thus the first column of the corresponding matrix for $S_{(j, (\alpha_1, \ldots, \alpha_p | \alpha_1, \ldots, \alpha_p))}$ is

$$\begin{aligned} (h_j, h_{\alpha_1}, \ldots, h_{\alpha_p}, & \overbrace{h_{-1}, \ldots, h_{-\alpha_p}}^{\alpha_p}, \overbrace{h_{-\alpha_p - 2}, \ldots, h_{-\alpha_{p-1}}}^{\alpha_{p-1} - \alpha_p - 1}, \ldots, \\ & \overbrace{h_{-\alpha_2 - 2}, \ldots, h_{-\alpha_1}}^{\alpha_1 - \alpha_2 - 1}, \overbrace{h_{-\alpha_1 - 2}, \ldots, h_{-m+1}}^{m - \alpha_1 - 2}). \end{aligned} \tag{3.40}$$

It is clearly that $S_{(j,(\alpha_1,\ldots,\alpha_p|\alpha_1,\ldots,\alpha_p))} \neq 0$ if and only if $\alpha_s \in \{0,\ldots,m-2\}\backslash\{j\}$ for $0 \leq j \leq m-2$, where $s \in \{1,\ldots,p\}$. In this case, there exists s s.t. $\alpha_{s+1} < j$ and $\alpha_s > j$ for $s \in \{1,\ldots,p\}$. Hence

$$\alpha_{s+1} + 2 \leq j+1, \qquad \alpha_s \geq j+1. \tag{3.41}$$

Since $(|(\alpha_1,\ldots,\alpha_p \mid \alpha_1,\ldots,\alpha_p)| + p)/2 = p + \alpha_1 + \cdots + \alpha_p$, thus

$$\begin{aligned} & (-1)^{p+\alpha_1+\cdots+\alpha_p} S_{(j,(\alpha_1,\ldots,\alpha_p|\alpha_1,\ldots,\alpha_p))} \\ = \; & (-1)^{p+\alpha_1+\cdots+\alpha_p+j+1} S_{(-(j+1),(\alpha_1,\ldots,\alpha_s,j,\alpha_{s+1},\ldots,\alpha_p|\alpha_1,\ldots,\alpha_s,j,\alpha_{s+1},\ldots,\alpha_p))}. \end{aligned} \tag{3.42}$$

We see that $S_{(-(j+1),(\alpha_1,\ldots,\alpha_p|\alpha_1,\ldots,\alpha_p))} \neq 0$ if and only if there exist $s \in \{\alpha_1,\ldots,\alpha_p\}$ s.t. $\alpha_s = j$ by (3.41), which implies that $A_j = A_{-(j+1)}$ for $j = 0,1,\ldots,m-2$. These complete the proof. □

In fact, Lemma 3.3 gives the character of the trivial module L_0 in form of an infinite sum of characters of generalized Verma modules.

Lemma 3.4. *For $\lambda \in \mathcal{P}$ and $\lambda_m = \lambda_0 = 0$, we have*

$$C_\lambda \cdot C_{\delta_1} = \sum_{\mu \in \mathcal{P}_\lambda} C_\mu. \tag{3.43}$$

Proof. Denote $T = S_{(1)} + S_{(0,\ldots,0,-1)} + y + 1 + y^{-1}$. We want to show that

$$C_\lambda \cdot T = \sum_{\mu \in \mathcal{P}_\lambda} C_\mu. \tag{3.44}$$

Since $C_0 = 1$ by Lemma 3.3, we get that $T = C_{\delta_1}$.

For convenience, we will write

$$A = \frac{\prod_{i=1}^{m}(1+x_i y) \prod_{i=1}^{m}(1+x_i y^{-1})}{\prod_{i=1}^{m}(1-x_i) \prod_{1\leq i<j\leq m}(1-x_i x_j)}. \tag{3.45}$$

Thus

$$\begin{aligned} & C_\lambda \cdot T \\ = \; & A\sum_{j=0}^{\infty}\sum_{\mu}\sum_{l=2}^{m}(-1)^{j+T(\mu)} S_{(j,\mu)\pm\delta_l}\sum_{s=0}^{2j} y^{j-s} + A\sum_{j=0}^{\infty}\sum_{\mu}(-1)^{j+T(\mu)} S_{(j,\mu)}\sum_{s=0}^{2j} y^{j-s} \\ + \; & A\sum_{\mu}(-1)^{T(\mu)} S_{(-1,\mu)} - A\sum_{\mu}(-1)^{T(\mu)} S_{(0,\mu)}. \end{aligned} \tag{3.46}$$

One can easily show that $\sum_{j=1}^{m} S_{\lambda\pm\delta_j} = 0$ if $S_\lambda = 0$. Moreover, $\sum_{j=1}^{m} S_{\lambda\pm\delta_j} = \sum_{j=1}^{m} S_{\lambda\pm\sigma(\delta_j)}$ for any $\sigma \in S_m \ltimes \mathbb{Z}_2^m$.

Using (3.19), we get that

$$\begin{aligned}(3.47)\qquad & C_\lambda \cdot T \\ = {} & A\sum_{j=0}^{\infty}\sum_{\sigma\in(S_{m-1}\ltimes\mathbb{Z}_2^{m-1})^\lambda}\sum_{l=2}^{m}(-1)^{j+l(\sigma)}S_{\rho^c-\sigma(\lambda^c+\rho^c\pm\delta_l)+j\delta_1}\sum_{s=0}^{2j}y^{j-s} \\ + {} & A\sum_{j=0}^{\infty}\sum_{\sigma\in(S_{m-1}\ltimes\mathbb{Z}_2^{m-1})^\lambda}(-1)^{j+l(\sigma)}S_{\rho^c-\sigma(\lambda^c+\rho^c)+j\delta_1}\sum_{s=0}^{2j}y^{j-s} \\ + {} & A\sum_{\sigma\in(S_{m-1}\ltimes\mathbb{Z}_2^{m-1})^\lambda}(-1)^{l(\sigma)}(S_{\rho^c-\sigma(\lambda^c+\rho^c)-\delta_1}-S_{\rho^c-\sigma(\lambda^c+\rho^c)}).\end{aligned}$$

We say that $\lambda^c\in\mathcal{P}^c_{\mathfrak{gl}(m)}$ if $\lambda\in\mathcal{P}_{\mathfrak{gl}(m)}$ (note that $\lambda_0=0$).

If $\lambda^c+\delta_{l+1}\notin\mathcal{P}^c_{\mathfrak{gl}(m)}$, i.e. $\lambda+\delta_{m-l}\notin\mathcal{P}_{\mathfrak{gl}(m)}$ for $l=1,\ldots,m-2$, then $\lambda_{m-l}=\lambda_{m-(l+1)}$.

Let $l\in\{j_1,\ldots,j_p\}$, for example, $l=j_{p-t}$. Now if $l+1\in\{j_1,\ldots,j_p\}$, then $l+1=j_{p-t+1}$. Thus $j_{p-t}+t+\lambda_{m-j_{p-t}}=j_{p-t+1}+t-1+\lambda_{m-j_{p-t+1}}$. It implies that $S_{\sigma(\lambda+\rho+\delta_{m-l})-\rho-j\delta_m}=0$. If $l+1\in\{r_1,\ldots,r_{m-1-p}\}$, we set $j'_k=j_k$ if $k\neq p-t$, and $j'_{p-t}=l+1$. Thus there exists $s\in\{1,\ldots,m-1-p\}$ s.t. $r_s=l+1$. We set $r'_q=r_q$ if $q\neq s$, and $r'_s=l$. We denote

$$w=(m-p-s,m-p+t)(0,1,\ldots,1,\overset{m-p-s}{-1},1,\ldots,1,\overset{m-p+t}{-1},1,\ldots,1). \tag{3.48}$$

Then $\sigma'=w\sigma$(note that $\sigma'\in\Gamma_{m-1}$), and

$$\sigma(\lambda+\rho+\delta_{m-l})-\rho-j\delta_m=\sigma'(\lambda+\rho+\delta_{m-l})-\rho-j\delta_m. \tag{3.49}$$

But $(-1)^{l(\sigma')}=-(-1)^{l(\sigma)}$, thus we know that

$$(-1)^{j+l(\sigma)}S_{\rho^c-\sigma(\lambda^c+\rho^c+\delta_{l+1})+j\delta_1}+(-1)^{j+l(\sigma')}S_{\rho^c-\sigma'(\lambda^c+\rho^c+\delta_{l+1})+j\delta_1}=0 \tag{3.50}$$

in (3.47). The case $l\in\{r_1,\ldots,r_{m-1-p}\}$ is similar.

If $\lambda^c-\delta_{l+2}\notin\mathcal{P}^c_{\mathfrak{gl}(m)}$, i.e. $\lambda-\delta_{m-l-1}\notin\mathcal{P}_{\mathfrak{gl}(m)}$ for $l=1,\ldots,m-2$, the discussion is similar to the above.

If $\lambda_{m-1}=0$, then $\lambda-\delta_{m-1}\notin\mathcal{P}_{\mathfrak{gl}(m)}$, i.e. $\lambda^c-\delta_2\notin\mathcal{P}^c_{\mathfrak{gl}(m)}$. Note that $\sigma(\lambda+\rho-\delta_{m-1})=\sigma(1,,\ldots,1,-1,1)(\lambda+\rho)$, and if $\sigma\in\Gamma_{m-1}$, then $\sigma'\in\Gamma_{m-1}$ by $\lambda_{m-1}=0$ and $\rho_{m-1}=\frac{1}{2}$. These imply that

$$\sum_{\sigma\in\Gamma_{m-1}}(-1)^{j+l(\sigma)}S_{\rho^c-\sigma(\lambda^c+\rho^c-\delta_2)+j\delta_1}+\sum_{\sigma\in\Gamma_{m-1}}(-1)^{j+l(\sigma)}S_{\rho-\sigma(\lambda^c+\rho^c)+j\delta_1}=0. \tag{3.51}$$

One shows that$\sum_\mu(-1)^{j_1+\cdots+j_p}S_{(-1,\mu)}-\sum_\mu(-1)^{j_1+\cdots+j_p}S_{(0,\mu)}=0$ if $\lambda_{m-1}=0$, via $S_{(0,\mu)}\neq0$ iff $j_1\neq1$ and $S_{(-1,\mu)}\neq0$ iff $j_1=1$. Thus (3.43) hold. $\square$

Proof of Theorem 3.1: We will prove the case $\lambda_0=0$ in this theorem by induction on the height of weight λ. Firstly, if $ht\lambda=0$ (i.e. $\lambda=0$), one easily shows that the statement holds by Lemma 3.2 and Lemma 3.3. Moreover, the statement holds for $\lambda=\delta_1$ by Lemma 3.2. Now by induction,

$$L_\lambda\otimes L_{\delta_1}=\sum_{\mu\in\mathcal{P}_\lambda}C_\mu\quad\text{if }\lambda_m=0. \tag{3.52}$$

In this case, λ is a tail atypical integral dominant weight. Since the weights in $\mathcal{P}_\lambda$ are in different blocks and $\tau_i\sigma(\lambda+\rho)-\rho-j\delta_i<\lambda$, then we have $L_\mu=C_\mu$ if $\mu\in\mathcal{P}_\lambda$ and if μ is also a tail atypical integral dominant weight by Remark 2.5. Now if $\lambda_{m-1}>0$, we consider $L_{\lambda+\delta_m}$. By (3.52) and above discussion, we can assume that

$$L_{\lambda+\delta_m}+xL_\lambda=L_\lambda+\sum_{\sigma\in\Gamma_m}(-1)^{\ell(\sigma)}M_{\sigma(\lambda+\delta_m+\rho)-\rho}, \tag{3.53}$$

where x is a nonnegative integer. One compares the coefficients of e^λ of the two sides of (3.53) then gets that $x<2$. But if $x=1$, one has $L_{\lambda+\delta_m}=\sum\limits_{\sigma\in\Gamma_m}(-1)^{\ell(\sigma)}M_{\sigma(\lambda+\delta_m+\rho)-\rho}$. It contradicts to Lemma 2.5. These prove the statement of the case $\lambda_0=0$.

Now we show *ii*) of the theorem with the case $\lambda_0>0$ (i.e. $k<m$ in (3.11)). Denote $\lambda=\sum_{j=1}^{k-1}\lambda_j\delta_j+\sum_{j=k}^{m}\delta_j+(m-k)\epsilon$ and $\lambda'=\sum_{j=1}^{k-1}\lambda_j\delta_j+\sum_{j=k}^{m}\delta_j+(m-k+1)\epsilon$. Then we claim that

$$F_{\lambda'}\cdot L_{\delta_1}=\sum_{\mu\in\mathcal{P}_{\lambda'}\backslash\{\lambda'\}}F_\mu. \tag{3.54}$$

This claim can be show by induction on k. Note that

$$\text{Left}=\sum_{\nu\in\mathcal{P}_{\lambda^T}}L_\nu+\sum_{\sigma\in\Gamma_m}\sum_{\xi\in\mathcal{P}^{(0)}_{\sigma(\lambda'+\rho)-\rho}}(-1)^{l(\sigma)}M_\xi, \tag{3.55}$$

$$\text{Right}=\sum_{\nu\in\mathcal{P}_{\lambda^T}}L_\nu+\sum_{\sigma\in\Gamma_m}\sum_{\eta\in\mathcal{P}_{\sigma(\lambda'+\rho)-\rho}\backslash\{\sigma(\lambda'+\rho)-\rho\}}(-1)^{l(\sigma)}M_\eta. \tag{3.56}$$

The elements in $\mathcal{P}^{(0)}_{\sigma(\lambda'+\rho)-\rho}$ but not in $\mathcal{P}_{\sigma(\lambda'+\rho)-\rho}\backslash\{\sigma(\lambda'+\rho)-\rho\}$ are $\nu=\sigma(\sum_{j=1}^{k-1}\lambda_j\delta_j+\sum_{j=k}^{m-1}\delta_j+(m-k+1)\epsilon+\rho)-\rho$ and $\zeta=\sigma(\sum_{j=1}^{k-1}\lambda_j\delta_j+\sum_{j=k}^{m}\delta_j+(m-k+1)\epsilon+\rho)-\rho$. Since $\sigma\cdot(1,\dots,1,-1)(\nu)=\zeta$ and $\sigma\cdot(1,\dots,1,-1)\in\Gamma_m$ if $\sigma\in\Gamma_m$. This implies that $\sum_{\sigma\in\Gamma_m}(-1)^{l(\sigma)}M_\nu+\sum_{\sigma\in\Gamma_m}(-1)^{l(\sigma)}M_\zeta=0$. Thus the claim is hold.

Now we show $L_\lambda=F_\lambda$ by induction on $m-k$. By (3.54), one has

$$L_\lambda=(1-x)L_{\lambda^T}+\sum_{\sigma\in\Gamma_m}(-1)^{l(\sigma)}M_{\sigma(\lambda+\rho)-\rho}, \tag{3.57}$$

where $x\geq 0$.

If $1<k<m$, the coefficient of $M_{\lambda^T+\delta_k}$ in $L_{\lambda'}$ or $L_{\lambda'^T}$ is nonnegative and in $\sum_{\sigma\in\Gamma_m}(-1)^{l(\sigma)}M_{\sigma(\lambda+\rho)-\rho}\otimes L_{\delta_1}$ is 0. One can tensor L_{δ_1} on both sides of (3.57) to show that $x\leq 1$. Lemma 2.5 says that $x=0$.

For $k=1$, note that $[L_{\sum_{j=1}^m\delta_j+(m-1)\epsilon}\otimes L_{\delta_1}:L_{\sum_{j=1}^m\delta_j+(m-2)\epsilon}]=1$. Denote $y=[L_{\sum_{j=1}^m\delta_j+(m-1)\epsilon}\otimes L_{\delta_1}:L_{\delta_1}]\geq 0$. On the other hand,

$$L_{\sum_{j=1}^m\delta_j+(m-1)\epsilon}\otimes L_{\delta_1}=(1-x)L_0\otimes L_{\delta_1}+\sum_{\sigma\in\Gamma_m}(-1)^{l(\sigma)}M_{\sigma(\sum_{j=1}^m\delta_j+(m-1)\epsilon+\rho)-\rho}\otimes L_{\delta_1}. \tag{3.58}$$

Note that by induction,

$$L_{\sum_{j=1}^m\delta_j+(m-2)\epsilon}=L_{\delta_1}+\sum_{\sigma\in\Gamma_m}(-1)^{l(\sigma)}M_{\sigma(\sum_{j=1}^m\delta_j+(m-2)\epsilon+\rho)-\rho}. \tag{3.59}$$

One substitutes (3.59) into (3.58) to show that $y=-x=0$. These complete the proof. □

4. Structure of tensor module $L_\lambda \otimes L_{\delta_1}$

Lemma 4.1. *For any $\mu \in \mathcal{P}_{\lambda^+}$, it should be that*

$$[L_\lambda \otimes L_{\delta_1} : L_\mu] = 1. \tag{4.1}$$

Particularly, if μ is typical, then L_μ is a direct summand in $L_\lambda \otimes L_{\delta_1}$.

Proof. If λ is a tail atypical weight, the statement has been shown in the proof of Theorem 3.1. Now assume λ is either typical or $(\delta_t + \epsilon)$-atypical $(1 \leq t \leq m)$. Thus $\chi_\lambda \neq \chi_{\lambda-\alpha}$ for any $\alpha \in \{\delta_j - \delta_k, \delta_l - \epsilon \mid 1 \leq j < k \leq m, 1 \leq l \leq m\}$. Hence for any $\nu \prec \lambda$ (i.e. $\chi_\nu = \chi_\lambda$ and $\nu < \lambda$), it should be that

$$\{\nu \pm \delta, \nu \pm \epsilon_j \mid 1 \leq j \leq m\} \cap \mathcal{P}_{\lambda^+} = \emptyset. \tag{4.2}$$

Therefore if we multiply L_{δ_1} on the both sides of (2.31) and calculate the right side by (3.6), then we can obtain that the coefficient of M_μ $(\mu \in \mathcal{P}_{\lambda^+})$ is exactly 1. Thanks to Remark 2.6, we get

$$[L_\lambda \otimes L_{\delta_1} : L_\mu] = 1 \quad \text{for any } \mu \in \mathcal{P}_{\lambda^+}. \tag{4.3}$$

□

Lemma 4.2. *For any $\lambda \in \mathcal{P}$, if L_μ is an irreducible submodule or quotient module of $L_\lambda \otimes L_{\delta_1}$, then it must be that $\mu \in \mathcal{P}_\lambda$.*

Proof. If λ is a tail atypical weight, then one can check that the statement hold by Theorem 3.1.

Now suppose that λ is either typical or $(\delta_t + \epsilon)$-atypical $(1 \leq t \leq m)$. Thus for any $\nu \prec \lambda$, there is no weight $\mu \in \mathcal{P}^0_\nu$ such that $\text{ht } \mu > \text{ht } \lambda$. Hence if we multiply L_{δ_1} on the right side of (2.31), then the coefficient of M_μ, where μ satisfies $\text{ht } \mu > \text{ht } \lambda$, is nonzero if and only if $\mu \in \mathcal{P}_{\lambda^+}$. So by Remark 2.6, we have

$$[L_\lambda \otimes L_{\delta_1} : L_\mu] = 0 \quad \text{for any } \mu \in \mathcal{P} \setminus \mathcal{P}_\lambda \text{ with ht } \mu > \text{ht } \lambda. \tag{4.4}$$

Take any irreducible submodule or quotient module L_μ of $L_\lambda \otimes L_{\delta_1}$.

Suppose $\text{ht } \mu > \text{ht } \lambda$. It must be that $\mu \in \mathcal{P}_\lambda$ because of (4.4).

Suppose $\text{ht } \mu < \text{ht } \lambda$. Since

$$\text{Hom}_{\mathfrak{g}}(L_\lambda \otimes L_{\delta_1}, L_\mu) \simeq \text{Hom}_{\mathfrak{g}}(L_\lambda, L_\mu \otimes L_{\delta_1}) \tag{4.5}$$

and

$$\text{Hom}_{\mathfrak{g}}(L_\mu, L_\lambda \otimes L_{\delta_1}) \simeq \text{Hom}_{\mathfrak{g}}(L_\mu \otimes L_{\delta_1}, L_\lambda), \tag{4.6}$$

it should be that L_λ is an irreducible submodule or quotient module of $L_\mu \otimes L_{\delta_1}$. Thus also by (4.4), we have $\lambda \in \mathcal{P}_\mu$, which implies that $\mu \in \mathcal{P}_\lambda$.

Suppose $\text{ht } \mu = \text{ht } \lambda$. There can not be a weight $\nu \succ \mu$ (note that $\text{ht } \nu - 2 \geq \text{ht } \mu = \text{ht } \lambda$) such that $[L_\lambda \otimes L_{\delta_1} : L_\nu] \neq 0$. Thus there should be a weight vector with highest weight μ. So if we multiply L_{δ_1} on the both sides of (2.31), then on the right side the coefficient of M_μ is nonzero. But it is clear by (3.6) that, for any $\nu \prec \lambda$ (note that $\text{ht } \nu \leq \text{ht } \lambda - 2$), the coefficient of M_μ in $M_\nu \otimes L_{\delta_1}$ is zero. So M_μ appears in $M_\lambda \otimes L_{\delta_1}$. It implies that $\mu \in \mathcal{P}_\lambda$. □

Lemma 4.3. *Suppose $\lambda \in \mathcal{P}$ is an atypical weight. For any $\mu, \nu \in \mathcal{P}_\lambda$ with $\mu \neq \nu$, it must be that $\chi_\mu \neq \chi_\nu$.*

Proof. Take any $\lambda \in \mathcal{P}$. If $\lambda_{m-1} = 0$ (which also implies $\lambda_m = \lambda_0 = 0$ by Theorem 2.1), one can check easily that the statement holds.

Now suppose that $\lambda_{m-1} \neq 0$, then $|\widetilde{\lambda}_1|, \ldots, |\widetilde{\lambda}_m|$ are pairwise distinguished.

If there exist $\mu \neq \nu \in \mathcal{P}_\lambda$ such that $\chi_\mu = \chi_\nu$. By Lemma 2.3, we only need to consider the case that μ and ν are both atypical. It is obvious by Lemma 2.3 that $\lambda - \mu, \lambda - \nu \in \{\pm\epsilon, 0\}$. Hence $|\widetilde{\mu}_0| = |\widetilde{\nu}_0|$. If either $\lambda - \mu$ or $\lambda - \nu$ is 0, then $\widetilde{\lambda_0} > 1$. Thus $|\widetilde{(\lambda \pm \epsilon)}_0| \neq |\widetilde{\lambda}_0|$, which is a contradiction to $|\widetilde{\mu}_0| = |\widetilde{\nu}_0|$. So $\mu, \nu = \lambda \pm \epsilon$, which is also a contradiction to $|\widetilde{\mu}_0| = |\widetilde{\nu}_0|$ since $|\widetilde{\mu}_0 - \widetilde{\nu}_0| = 2$ and $\widetilde{\nu}_0, \widetilde{\mu}_0 \in \mathbb{Z} + \frac{1}{2}$. □

Corollary 4.4. *For any atypical weight $\lambda \in \mathcal{P}$, if $\mu \in \mathcal{P}_\lambda$ is also an atypical weight, then L_μ is a direct summand in $L_\lambda \otimes L_\delta$ and*

$$[L_\lambda \otimes L_\delta : L_\mu] = 1. \tag{4.7}$$

Proof. Just combine Lemmas 4.1, 4.2 and 4.3. □

Remark 4.5. Lemma 4.3 and Corollary 4.4 indicate that for any two atypical weights $\lambda, \mu \in \mathcal{P}$, if there exist atypical weights $\mu^{(0)} = \lambda, \mu^{(1)}, \cdots, \mu^{(t)} = \mu \in \mathcal{P}$ such that $\mu^{(i+1)} \in \mathcal{P}_{\mu^{(i)}}$, then one can use (3.7) iteratively to get ch L_μ from ch L_λ by a straightforward calculation:

$$L_{\mu^{(i)}} \xrightarrow{\otimes L_\delta} L_{\mu^{(i+1)}}. \tag{4.8}$$

Such an algorithm will be used in the next section.

Remark 4.6. All of the results stated in this section are also valid for $\mathfrak{osp}(n|2)$ (cf. [9]).

5. Character formulae for non-tail atypical weights

5.1. When $\varphi(\lambda) = \lambda^T$.

We have dealt with this case in Theorem 3.1 ii).

5.2. When $\varphi(\lambda) \neq \lambda^T$.

5.2.1. **In case that λ is $(\delta_m + \epsilon)$-atypical.** In this case, $\lambda_m = \lambda_0 + 1 > 1$ and $\lambda^T = \lambda - \lambda_0(\delta_m + \epsilon) - \delta_m$. Assume that $\lambda_0 = s > 0$.

Let $\mu = \lambda - \epsilon$. It is obvious that $\mu \in \mathcal{P}$ is typical. So by Lemma 2.5, we have

$$L_\mu = \sum_{\sigma \in \Gamma_m} (-1)^{\ell(\sigma)} M_{\sigma(\mu+\rho)-\rho}. \tag{5.1}$$

Suppose

$$[L_\mu \otimes L_{\delta_1} : L_{\lambda - i(\delta_m+\epsilon)}] = x_i, (0 \leq i \leq s); \quad [L_\mu \otimes L_{\delta_1} : L_{\lambda - s(\delta_m+\epsilon)-\delta_m}] = y. \tag{5.2}$$

Note that $x_0 = 1$ by Lemma 4.1.

Multiply L_{δ_1} on the both sides of the above formula. By (3.7), we can calculate that the right side becomes to

$$\sum_{\nu \in \mathcal{P}_\mu} \sum_{\sigma \in \Gamma_m} (-1)^{\ell(\sigma)} M_{\sigma(\nu+\rho)-\rho}. \tag{5.3}$$

Select in (5.3) the modules M_ν with $\chi_\nu = \chi_\lambda$. Then we have

$$L_\lambda + \sum_{i=1}^{s} x_i L_{\lambda - i(\delta_m+\epsilon)} + y L_{\lambda - s(\delta_m+\epsilon)-\delta_m} = \sum_{\sigma \in \Gamma_m} (-1)^{\ell(\sigma)} (M_{\sigma(\lambda+\rho)-\rho} + M_{\sigma(\lambda-\delta_m-\epsilon+\rho)-\rho}). \tag{5.4}$$

By Lemma 4.1, we have that

$$[L_{\lambda-\delta_m-\epsilon}\otimes L_{\delta_1}:L_\mu]=1. \tag{5.5}$$

Furthermore, since μ is typical, we can get from Lemma 4.2 that

$$[L_{\lambda-i(\delta_m+\epsilon)}\otimes L_{\delta_1}:L_\mu]=[L_{\lambda-s(\delta_m+\epsilon)-\delta_m}\otimes L_{\delta_1}:L_\mu]=0,\quad (i=2,3,\ldots,s). \tag{5.6}$$

On the other hand, also thanks to that μ is typical, it is not difficult to calculate that

$$[\sum_{\sigma\in\Gamma_m}(-1)^{\ell(\sigma)}(M_{\sigma(\lambda+\rho)-\rho}+M_{\sigma(\lambda-\delta_m-\epsilon+\rho)-\rho})\otimes L_{\delta_1}:L_\mu]=2 \tag{5.7}$$

Combining (5.4)-(5.7), we get that

$$x_1\le[L_\lambda\otimes L_{\delta_1}:L_\mu]+x_1=2. \tag{5.8}$$

Moreover, since

$$\dim\operatorname{Hom}_{\mathfrak g}(L_\mu\otimes L_{\delta_1},L_{\mu-\delta_m})=\dim\operatorname{Hom}_{\mathfrak g}(L_\mu,L_{\mu-\delta_m}\otimes L_{\delta_1})=1, \tag{5.9}$$

it should be that $x_1>0$.

Below we shall use induction on s to obtain L_λ.

1). For s=1. Then by Theorem 3.1 ii), one can rewrite (5.4) as

$$L_\lambda+x_1L_{\lambda-\delta_m-\epsilon}+yL_{\lambda-2\delta_m-\epsilon}=\sum_{\sigma\in\Gamma_m}(-1)^{\ell(\sigma)}M_{\sigma(\lambda+\rho)-\rho}+L_{\lambda-\delta_m-\epsilon}-L_{\lambda-2\delta_m-\epsilon}. \tag{5.10}$$

That is,

$$L_\lambda=\sum_{\sigma\in\Gamma_m}(-1)^{\ell(\sigma)}M_{\sigma(\lambda+\rho)-\rho}+(1-x_1)L_{\lambda-\delta_m-\epsilon}-(1+y)L_{\lambda-2\delta_m-\epsilon}. \tag{5.11}$$

Multiply L_{δ_1} on the both sides of the above equation and select the terms L_ν and M_ν with $\chi_\nu=\chi_\lambda$, then we get

$$L_\lambda=\sum_{\sigma\in\Gamma_m}(-1)^{\ell(\sigma)}M_{\sigma(\lambda+\rho)-\rho}+(1-x_1)L_{\lambda-2\delta_m-\epsilon}-(1+y)L_{\lambda-\delta_m-\epsilon}. \tag{5.12}$$

Comparing (5.11) with (5.12) shows us that

$$x_1-1=y+1\quad\Rightarrow\quad x_1=2+y\ge 2. \tag{5.13}$$

Inequalities (5.8) and (5.13) induce that $x_1=2$ and $y=0$. Thus

$$L_\lambda=\sum_{\sigma\in\Gamma_m}(-1)^{\ell(\sigma)}M_{\sigma(\lambda+\rho)-\rho}-L_{\lambda-\delta_m-\epsilon}-L_{\lambda-2\delta_m-\epsilon}. \tag{5.14}$$

As a co-product,

$$[L_\lambda\otimes L_{\delta_1}:L_{\lambda-\delta_m}]=[L_\lambda\otimes L_{\delta_1}:L_{\lambda-\epsilon}]=0. \tag{5.15}$$

2). For s=2. Now using (5.14), equation (5.4) can be rewritten as

$$\begin{aligned}&L_\lambda+x_1L_{\lambda-\delta_m-\epsilon}+x_2L_{\lambda-2\delta_m-2\epsilon}+yL_{\lambda-3\delta_m-2\epsilon}\\&=\sum_{\sigma\in\Gamma_m}(-1)^{\ell(\sigma)}M_{\sigma(\lambda+\rho)-\rho}+L_{\lambda-\delta_m-\epsilon}+L_{\lambda-2\delta_m-2\epsilon}+L_{\lambda-3\delta_m-2\epsilon}.\end{aligned} \tag{5.16}$$

That is,

$$L_\lambda=\sum_{\sigma\in\Gamma_m}(-1)^{\ell(\sigma)}M_{\sigma(\lambda+\rho)-\rho}+(1-x_1)L_{\lambda-\delta_m-\epsilon}+(1-x_2)L_{\lambda-2\delta_m-2\epsilon}+(1-y)L_{\lambda-3\delta_m-2\epsilon}. \tag{5.17}$$

Notice that $\lambda-\delta_m-2\epsilon$ is a typical weight, hence

$$[L_\lambda\otimes L_{\delta_1}:L_{\lambda-\delta_m-2\epsilon}]=[L_{(\lambda-2\delta_m-2\epsilon)^\delta}\otimes L_{\delta_1}:L_{\lambda-\delta_m-2\epsilon}]=0 \tag{5.18}$$

by Lemma 4.2,

$$[\sum_{\sigma\in\Gamma_m}(-1)^{\ell(\sigma)}M_{\sigma(\lambda+\rho)-\rho}\otimes L_{\delta_1}:L_{\lambda-\delta-2\epsilon_k}]=0 \tag{5.19}$$

by direct calculation,

$$[L_{\lambda-\delta_m-\epsilon}\otimes L_{\delta_1}:L_{\lambda-\delta_m-2\epsilon}]=0 \tag{5.20}$$

by (5.15), and

$$[L_{\lambda-2\delta_m-2\epsilon}\otimes L_{\delta_1}:L_{\lambda-\delta_m-2\epsilon}]=1 \tag{5.21}$$

by Lemma 4.1.

Therefore (5.17)-(5.21) imply that $x_2=1$.

Multiply L_δ on the both sides of (5.17) and select the terms L_ν and M_ν with $\chi_\nu=\chi_\lambda$, then we have

$$L_\lambda=\sum_{\sigma\in\Gamma_m}(-1)^{\ell(\sigma)}M_{\sigma(\lambda+\rho)-\rho}+(1-x_1)L_{\lambda-\delta_m-\epsilon}+(1-x_2)L_{\lambda-3\delta_m-2\epsilon}+(1-y)L_{\lambda-2\delta_m-2\epsilon}. \tag{5.22}$$

Compare (5.17) and (5.22), then we get

$$1-x_2=1-y \quad\Rightarrow\quad y=x_2=1. \tag{5.23}$$

Therefore,

$$L_\lambda=\sum_{\sigma\in\Gamma_m}(-1)^{\ell(\sigma)}M_{\sigma(\lambda+\rho)-\rho}+(1-x_1)L_{\lambda-\delta_m-\epsilon}. \tag{5.24}$$

By (5.8) and (5.9), we know that $x_1=1$ or 2. But it is impossible that $x_1=1$ because of Lemma 2.5. So $x_1=2$ and

$$L_\lambda=\sum_{\sigma\in\Gamma_m}(-1)^{\ell(\sigma)}M_{\sigma(\lambda+\rho)-\rho}-L_{\lambda-\delta_m-\epsilon}. \tag{5.25}$$

Now it is easy to get from the above equation that

$$[L_\lambda\otimes L_{\delta_1}:L_{\lambda-\delta_m}]=[L_\lambda\otimes L_{\delta_1}:L_{\lambda-\epsilon}]=0. \tag{5.26}$$

3). For any s $\geq$ 2. We shall use induction on s to show that (5.25) and (5.26) hold for any $s\geq 2$. By induction assumption, we can get from (5.4) that

$$L_\lambda=\sum_{\sigma\in\Gamma_m}(-1)^{\ell(\sigma)}M_{\sigma(\lambda+\rho)-\rho}+\sum_{i=1}^{2}(1-x_i)L_{\lambda-i\delta_m-i\epsilon}-\sum_{i=3}^{s}x_iL_{\lambda-i\delta_m-i\epsilon}-yL_{\lambda-(s+1)\delta_m-s\epsilon}. \tag{5.27}$$

Multiply L_{δ_1} on both sides of the above formula. Then if we choose the terms M_ν and L_ν with $\chi_\nu=\chi_{\lambda-i\delta_m-(i+1)\epsilon}(1\leq i<s)$, there comes that $x_2=1$, $x_3=\cdots=x_s=0$. Moreover, if we choose the terms M_ν and L_ν with $\chi_\nu=\chi_\lambda$, there comes that $y=x_s=0$.

Thus

$$L_\lambda=\sum_{\sigma\in\Gamma_m}(-1)^{\ell(\sigma)}M_{\sigma(\lambda+\rho)-\rho}+(1-x_1)L_{\lambda-\delta_m-\epsilon}. \tag{5.28}$$

Recall that we have known $x_1 = 1$ or 2. But only the case of $x_1 = 2$ is valid by (5.28) because of Lemma 2.5. Hence we have shown that (5.25) and (5.26) hold for any $s \geq 2$.

5.2.2. **In case that λ is $(\delta_k + \epsilon)$-atypical, $(k < m)$.** In this case, $\lambda_0 + 1 + k - m = \lambda_k > 1$ and $\lambda_k \geq \lambda_{k+1} \geq \lambda_{k+2} \geq \cdots \geq \lambda_m \geq 1$.

Let

$$\lambda^{(t)} = \sum_{i=1}^{k} \lambda_i \delta_i + \sum_{i=k+1}^{m} \lambda_k \delta_i + (\lambda_k - 1 + t)\epsilon, \quad (t = 0, 1 \ldots, m-k), \tag{5.29}$$

$$\lambda^{(m-k+t)} = \sum_{i=1}^{k} \lambda_i \delta_i + \sum_{i=k+1}^{m} \lambda_k \delta_i + \lambda_0 \epsilon - t\delta_m, \quad (t = 1, 2, \ldots, \lambda_k - \lambda_m),$$

$$\lambda^{(m-k+\lambda_k-\lambda_m+t)} = \sum_{i=1}^{k} \lambda_i \delta_i + \sum_{i=k+1}^{m-1} \lambda_k \delta_i + \lambda_m \delta_m + \lambda_0 \epsilon - t\delta_{m-1}, \quad (t = 1, 2, \ldots, \lambda_k - \lambda_{m-1}),$$

$$\cdots\cdots\cdots\cdots\cdots\cdots$$

$$\lambda^{(m-k+\sum_{i=k+2}^{m}(\lambda_k-\lambda_i)+t)} = \sum_{i=1}^{k} \lambda_i \delta_i + \lambda_k \delta_{k+1} + \sum_{i=k+1}^{m} \lambda_i \delta_i + \lambda_0 \epsilon - t\delta_{k+1}, \quad (t = 1, 2, \ldots, \lambda_k - \lambda_{k+1}),$$

$$\lambda^{(m-k+\sum_{i=k+1}^{m}(\lambda_k-\lambda_i))} = \lambda.$$

Notice that all $\lambda^{(t)}$ $(0 \leq t \leq m - k + \sum_{i=k+1}^{m}(\lambda_k - \lambda_i))$ are atypical weights and $\lambda^{(t+1)} \in \mathcal{P}_{\lambda^{(t)}}$.

We have obtained in Section 5.2.1 that

$$L_{\lambda^{(0)}} = \sum_{\sigma \in \Gamma_m} (-1)^{\ell(\sigma)} M_{\sigma(\lambda^{(0)}+\rho)-\rho} - L_{\varphi(\lambda^{(0)})}. \tag{5.30}$$

Using the algorithm introduced in Remark 4.5, now there is no difficulty for us to calculate that

$$L_{\lambda} = \sum_{\sigma \in \Gamma_m} (-1)^{\ell(\sigma)} M_{\sigma(\lambda+\rho)-\rho} - L_{\varphi(\lambda)} \tag{5.31}$$

by (5.29) and (5.30).

5.2.3. **Final results for the case $\varphi(\lambda) \neq \lambda^T$.**

To summarize:

Theorem 5.1. *For any non-tail atypical weight λ with $\varphi(\lambda) \neq \lambda^T$,*

$$L_{\lambda} = \begin{cases} \sum_{\sigma \in \Gamma_m} (-1)^{\ell(\sigma)} M_{\sigma(\lambda+\rho)-\rho} - L_{\varphi(\lambda)}, & \textit{if } \varphi^2(\lambda) \neq \lambda^T \\ \sum_{\sigma \in \Gamma_m} (-1)^{\ell(\sigma)} M_{\sigma(\lambda+\rho)-\rho} - L_{\varphi(\lambda)} - L_{\lambda^T}, & \textit{if } \varphi^2(\lambda) = \lambda^T. \end{cases} \tag{5.32}$$

For any atypical weight $\lambda \in \mathcal{P}$, denote by θ_λ the unique number such that $\varphi^{\theta_\lambda+1}(\lambda) = \lambda^T$.

Theorem 5.2. *For any atypical weight $\lambda \in \mathcal{P}$ with $\theta_\lambda \geq 1$, one has*

$$L_\lambda = \sum_{i=0}^{\theta_\lambda} \sum_{\sigma \in \Gamma_m} (-1)^{i+\ell(\sigma)} M_{\sigma(\varphi^i(\lambda)+\rho)-\rho} + \tag{5.33}$$

$$\sum_{\sigma \in \Gamma_{m-1}} \sum_{i=\flat(\sigma,\lambda^T)}^{m} \sum_{j=max\{0,\frac{1}{2}-\sigma(\lambda^T+\rho)_{i-1}\}}^{-\frac{3}{2}-\tau_i\sigma(\lambda^T+\rho)_{i+1}} 2(-1)^{\theta_\lambda+\ell(\tau_i\sigma)+j} M_{\tau_i\sigma(\lambda^T+\rho)-\rho+j\epsilon-j\delta_i}.$$

References

1. J. Brundan, *Kazhdan-Lusztig polynomials and character formulae for the Lie superalgebra* $\mathfrak{gl}(m|n)$, J. Amer. Math. Soc. **16** (2003), 185–231.
2. S.-J. Cheng, N. Lam and W. Wang, *Super duality and irreducible characters of ortho-symplectic Lie superalgebras*, arXiv: 0911.0129.
3. S.-J. Cheng, W. Wang and R.B. Zhang, *Super duality and Kazhdan-Lusztig polynomials*, Trans. Amer. Math. Soc. **360** (2008), 5883–5924.
4. S.-J. Cheng and R.B. Zhang, *Analogue of Kostant's* $\mathfrak{u}$*-cohomology formula for the general linear superalgebra*, Int. Math. Res. Not. (2004), 31–53.
5. C. Gruson and V. Serganova, *Cohomology of generalized supergrassmannians and character formulae for basic classical Lie superalgebras*, arXiv: 0906.0918.
6. V.G. Kac, *Lie superalgebras*, Adv. Math. **26** (1977), 8–96.
7. V.G. Kac, *Characters of typical representations of classical Lie superalgebras*, Comm. Algebra **5** (1977), 889–897.
8. V.G. Kac, *Representations of classical Lie superalgebras*, Lect. Notes Math. 676 (1978), 597–626.
9. L. Luo, *Character formulae for ortho-symplectic Lie superalgebras* $\mathfrak{osp}(n|2)$, arXiv: 0909.3134.
10. I.G. Macdonald, *Symmetric functions and Hall polynomials*, Second Edition, Clarendon Press, Oxford, 1995.
11. V. Serganova, *Kazhdan-Lusztig polynomials and character formula for the Lie superalgebra* $\mathfrak{gl}(m|n)$, Selecta. Math. (N.S.) **2** (1996), 607–651.
12. V. Serganova, *Characters of irreducible representations of simple Lie superalgebras*, Proceedings of the International Congress of Mathematicians 1998, Berlin, Vol. II, 583–593.
13. Y. Su and R.B. Zhang, *Character and dimension formulae for general linear superalgebra*, Adv. Math. **211** (2007), no. 1, 1–33.
14. Y. Su and R.B. Zhang, *Generalised Verma modules for the orthosymplectic Lie superalgebra* $\mathfrak{osp}_{k|2}$, arXiv:0911.0735.
15. J. van der Jeugt, *Irreducible representations of the exceptional Lie superalgebras* $D(2,1;\alpha)$, J. Math. Phys. **26** (1985), 913–924.
16. J. van der Jeugt, *Character formulae for the Lie superalgebra* $C(n)$, Comm. Algebra **19** (1991), 199–222.

Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing, 100190, China
E-mail address: `caobintao@amss.ac.cn`

Department of Mathematics, East China Normal University, Shanghai, 200241, China
E-mail address: `lluo@math.ecnu.edu.cn`